



Chilliwack FC Concussion Management Policy

POLICY OVERVIEW:

A concussion is a brain injury. All concussions should be regarded as potentially serious. Most concussions recover completely with correct management. Incorrect management of a concussion can lead to further injury. Concussions should be managed according to current guidelines. Anyone with suspected concussion following an injury must be immediately removed from playing or training and receive a prompt assessment by a medical doctor or nurse practitioner. Concussions are managed by licensed health care professionals working within their scope of practice and expertise. Concussions are managed by a limited period of rest followed by avoiding physical and brain activities that make concussive symptoms worse, and once concussion related symptoms have resolved, a stepwise return to school, work and sports-related activities. Return to education or work must take priority over return to playing soccer. Concussion symptoms must have completely resolved and documented medical clearance completed by a medical doctor or nurse practitioner must be received before resuming full contact practice or game play. The recurrence of concussion symptoms subsequent to the return to full contact practice or game play requires removal from training or playing and reassessment.

PRELIMINARY STATEMENT:

Chilliwack FC is adopting this policy as part of its obligation to provide a safe playing environment for its players.

Awareness of concussion and effects of concussion on players, especially youth players, has increased over the past decade. Medically, a concussion is a form of traumatic brain injury. Because it is a brain injury, a concussion is not a visible injury, such as a broken arm. It requires time to heal properly, involving both physical and mental rest. Consequences may include loss of coordination or memory, vision problems, learning issues and changes in emotions and behavior.

A concussion may result from a direct blow to the head or from an indirect blow to the body that causes various forces to affect the brain within the skull. Signs and symptoms may appear immediately or may be noticed hours or days (or longer) later. There are a wide variety of symptoms which are possible. Loss of consciousness is not required to be diagnosed with a concussion. Adolescents require more time to recover from concussions than adults.

APPLICATION OF POLICY:

This policy applies to all Chilliwack FC coaches, team managers, officials and players who are involved in Chilliwack FC programs.

REQUIREMENTS:

All Chilliwack FC coaches, team managers and club officials are required to take the free CDC (Center for Disease Control) online training course “Concussion Training for Coaches” before they may coach. This rule goes into effect with the start of the 2015 spring soccer season. The certificate of completion at the end of the online course is proof the course has been taken. This course takes approx. 35-40 mins to complete. Please submit all certificates to info@chilliwackfc.com

The training course can be accessed by clicking this link:

http://www.cdc.gov/concussion/HeadsUp/online_training.html

This document contains references to Soccer Canada and BC Soccer Concussion Policies



Chilliwack FC Concussion Management Policy

As part of a pre-season concussion education strategy, CDC (Center for Disease Control) online training course “Concussion Training for Coaches” However, the CDC offers additional training for non-coaches. The Chilliwack FC strongly recommends this training for all involved with the sport.

- Players/Coaches, and parents/guardians should review this document at a minimum regarding Concussion Management as part of their soccer registration process.
- All participants in our sport should be encouraged to familiarize themselves with the entirety of our Concussion Guidelines. Baseline (pre-season) testing of youth (<18 years) and adult recreational athletes using any tool or combination of tools is not required for post-injury care of those who sustain a suspected or diagnosed concussion and is not recommended. For the communication between physicians and soccer coaches, team officials and clubs we recommend using:

Canada Soccer’s Concussion Assessment Report, available @ canadasoccer.com.
Generic concussion reporting letters are also available through Parachute Canada:
http://www.parachutecanada.org/downloads/injurytopics/Medical-Assessment-Letter_Parachute.pdf & http://www.parachutecanada.org/downloads/injurytopics/Medical-Clearance-Letter_Parachute.pdf

Concussion Information from Soccer Canada

This policy is intended for those managing concussion in soccer at all levels. Professional and National level players typically have access to an enhanced level of medical care, which means that their concussion and their return to play can be managed in a more closely monitored way. This Chilliwack FC Policy is based on current evidence and examples of best practice taken from soccer organizations around the world and other sports, including Soccer Canada, the Football Association, the Scottish FA, World Rugby, and the Canadian Concussion Collaborative. They are consistent with The Canadian Guideline on Concussion in Sport, (Toronto: Parachute, 2017) and the current Consensus Statement on Concussion in Sport issued by the Fifth International Conference on Concussion in Sport, Berlin 2017. The Policy has been reviewed and is approved by Canada Soccer Sports Medicine Committee. While this policy aims to reflect ‘best practice’, it must be recognized that there is a current lack of evidence with respect to their effectiveness in preventing long-term harm.

Canada Soccer Sports Medicine Committee will continue to monitor research and consensus in the area of concussion and update its policies accordingly. This version was adopted by Canada Soccer in 2018 and should be reviewed no later than 2022. Chilliwack FC will have a practice of “best practices” as it relates to this policy based on current and evidence-based research and resources available for Concussion Management.

ADDITIONAL RESOURCES WEBSITES:

Sport Information Resource Centre (SIRC):

<http://sirc.ca/resources/concussion>

Coaching Association of Canada: Making Head Way Concussion eLearning Series:

<https://www.coach.ca/concussionawareness-s16361>

Parachute: www.parachutecanada.org/concussion

Concussion Awareness Training Tool: www.catonline.com Ontario Ministry of Health Concussion



Chilliwack FC Concussion Management Policy

Resources: <http://www.health.gov.on.ca/en/public/programs/concussions/>

VIDEOS:

Dr. Mike Evans Health Lab — Concussions: <https://www.reframehealthlab.com/concussions/>

BC Soccer references:

The British Columbia government has established a website with key partners, which is available at: <http://www.healthlinkbc.ca/healthtopics/content.asp?hwid=tp23364spec>

WE ALL NEED TO PLAY A PART IN THE RECOGNITION AND MANAGEMENT OF CONCUSSION

We have a heightened awareness of concussions, related to increased media coverage of this brain injury with its range of outcomes, incidents involving high profile athletes with concussion, and increasing understanding of the consequences of repetitive brain trauma, primarily within professional sports.

WHAT IS A “CONCUSSION”? Concussion is an injury to the brain resulting in a disturbance of brain function involving thinking and behavior.

WHAT CAUSES CONCUSSION? Concussion can be caused by a direct blow to the head or an impact to the body causing rapid movement of the head.

ONSET OF SYMPTOMS Symptoms of concussion typically appear immediately but may evolve within the first 24–48 hours.

WHO IS AT RISK? All our sport’s participants (players, but also team staff and officials). Some soccer participants are at increased risk of concussion: Children and adolescents (18 years and under) are more susceptible to brain injury, take longer to recover, and are susceptible to rare dangerous brain complications, which may include death. Female soccer players have higher rates of concussion. Participants with previous concussion are at increased risk of further concussions — which may take longer to recover.

WHAT ARE THE DANGERS OF BRAIN INJURY? Failure to recognize and report concussive symptoms or returning to activity with ongoing concussion symptoms set the stage for: 1. Cumulative concussive injury 2. ‘Second Impact Syndrome’ Second impact syndrome is a rare occurrence. An athlete sustains a brain injury and while still experiencing symptoms (not fully recovered), sustains a second brain injury, which is associated with brain swelling and permanent brain injury or death. Brain swelling may also occur without previous trauma. Recurrent brain injury is currently implicated in the development of Chronic Traumatic Encephalopathy Chronic Traumatic Encephalopathy (CTE) is a progressive degenerative brain disease seen in people with a history of brain trauma. For athletes, the brain trauma has been repetitive. Originally described in deceased boxers, it now has been recognized in many sports. Symptoms include difficulty thinking, explosive and aggressive behavior, mood disorder (depression), and movement disorder (parkinsonism).



Chilliwack FC Concussion Management Policy

RECOGNIZE — LEARN THE SIGNS AND SYMPTOMS OF A CONCUSSION SO YOU UNDERSTAND WHEN A SOCCER PLAYER MIGHT HAVE A SUSPECTED CONCUSSION.

Everyone involved in the game (including staff, coaches, officials, players, parents and guardians of children and adolescents) should be aware of the signs, symptoms, and dangers of concussion. If any of the following signs or symptoms are present following an injury the player should be suspected of having concussion and immediately removed from play or training.

- “If in doubt, sit them out.”
- “It is better to miss one game than the whole season.”

VISIBLE CLUES OF CONCUSSION — WHAT YOU MAY SEE: Any one or more of the following visual clues can indicate a concussion:

- Dazed, blank or vacant look
- Lying motionless on ground / slow to get up
- Unsteady on feet / balance problems / falling over / poor coordination
- Loss of consciousness or responsiveness
- Confused or not aware of play or events
- Grabbing, clutching, or shaking of the head
- Seizure
- More emotional or irritable than normal for that person
- Injury event that could have caused a concussion

SYMPTOMS OF CONCUSSION — WHAT YOU MAY BE TOLD BY AN INJURED PLAYER:

The presence of any one or more of the following symptoms may suggest a concussion:

- Headache or “Pressure in head”
- Dizziness or balance problems
- Mental clouding, confusion, or feeling slowed down
- Trouble seeing
- Nausea or vomiting
- Fatigue
- Drowsiness or feeling like “in a fog “or difficulty concentrating
- Sensitivity to light or noise
- Difficulty with reading, learning or work
- Sleep problems: getting asleep, too much or too little
- Emotional / anger / sad / anxious

The **Concussion Recognition Tool 5** is valuable for all first responders in recognizing suspected concussion and responding to more severe brain injury or potential neck injury.



Chilliwack FC Concussion Management Policy

CONCUSSION RECOGNITION TOOL 5 ©

To help identify concussion in children, adolescents and adults



Supported by



RECOGNISE & REMOVE

Head impacts can be associated with serious and potentially fatal brain injuries. The Concussion Recognition Tool 5 (CRT5) is to be used for the identification of suspected concussion. It is not designed to diagnose concussion.

STEP 1: RED FLAGS – CALL AN AMBULANCE

If there is concern after an injury including whether ANY of the following signs are observed or complaints are reported then the player should be safely and immediately removed from play/game/activity. If no licensed healthcare professional is available, call an ambulance for urgent medical assessment:

- Neck pain or tenderness
- Double vision
- Weakness or tingling/ burning in arms or legs
- Severe or increasing headache
- Seizure or convulsion
- Loss of consciousness
- Deteriorating conscious state
- Vomiting
- Increasingly restless, agitated or combative

Remember:

- In all cases, the basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Assessment for a spinal cord injury is critical.
- Do not attempt to move the player (other than required for airway support) unless trained to do so.
- Do not remove a helmet or any other equipment unless trained to do so safely.

If there are no Red Flags, identification of possible concussion should proceed to the following steps:

STEP 2: OBSERVABLE SIGNS

Visual clues that suggest possible concussion include:

- Lying motionless on the playing surface
- Slow to get up after a direct or indirect hit to the head
- Disorientation or confusion, or an inability to respond appropriately to questions
- Blank or vacant look
- Balance, gait difficulties, motor incoordination, stumbling, slow laboured movements
- Facial injury after head trauma

© Concussion in Sport Group 2017

STEP 3: SYMPTOMS

- Headache
- "Pressure in head"
- Balance problems
- Nausea or vomiting
- Drowsiness
- Dizziness
- Blurred vision
- Sensitivity to light
- Sensitivity to noise
- Fatigue or low energy
- "Don't feel right"
- More emotional
- More Irritable
- Sadness
- Nervous or anxious
- Neck Pain
- Difficulty concentrating
- Difficulty remembering
- Feeling slowed down
- Feeling like "in a fog"

STEP 4: MEMORY ASSESSMENT

(IN ATHLETES OLDER THAN 12 YEARS)

Failure to answer any of these questions (modified appropriately for each sport) correctly may suggest a concussion:

- "What venue are we at today?"
- "Which half is it now?"
- "Who scored last in this game?"
- "What team did you play last week/game?"
- "Did your team win the last game?"

Athletes with suspected concussion should:

- Not be left alone initially (at least for the first 1-2 hours).
- Not drink alcohol.
- Not use recreational/ prescription drugs.
- Not be sent home by themselves. They need to be with a responsible adult.
- Not drive a motor vehicle until cleared to do so by a healthcare professional.

The CRT5 may be freely copied in its current form for distribution to individuals, teams, groups and organisations. Any revision and any reproduction in a digital form requires approval by the Concussion in Sport Group. It should not be altered in any way, rebranded or sold for commercial gain.

ANY ATHLETE WITH A SUSPECTED CONCUSSION SHOULD BE IMMEDIATELY REMOVED FROM PRACTICE OR PLAY AND SHOULD NOT RETURN TO ACTIVITY UNTIL ASSESSED MEDICALLY, EVEN IF THE SYMPTOMS RESOLVE

© Concussion in Sport Group 2017



Chilliwack FC Concussion Management Policy

REMOVE - If a soccer player has a suspected concussion he or she must be removed from activity immediately. Team-mates, staff, coaches, players or parents and guardians who suspect that a player may have concussion **MUST** work together to ensure that the player is removed from play in a safe manner. If a neck injury is suspected the player should not be moved and should only be removed from the field of play by emergency healthcare professionals with appropriate spinal care training. Call 911. More severe forms of brain injury may be mistaken for concussion. If **ANY** of the following are observed or reported within 48 hours of an injury, then the player should be transported for urgent medical assessment at the nearest hospital (symptoms below).

Call 911. Activate your emergency action plan.

- Neck pain or tenderness
- Deteriorating consciousness (drowsier)
- Increasing confusion or irritability Severe or increasing headache
- Repeated vomiting
- Unusual behavior change
- Seizure Double vision
- Weakness or tingling / burning in arms or legs

Anyone with a suspected concussion should not: be left alone until they have been assessed medically consume alcohol or recreational drugs in the first 24 hours, and thereafter should avoid alcohol or recreational drugs until free of all concussion symptoms drive a motor vehicle until cleared to do so by a medical doctor or nurse practitioner

RE-ENTRY — A licensed healthcare professional with **expertise** in the evaluation and management of head injury and concussions may review a player with suspected concussion at field side. A player who has been removed from play who reports **NO** concussion symptoms and **NO** visual clues of a concussion can be returned to play. Any such player should be monitored for delayed symptoms, which may appear over the next 24-48 hours. If there is any doubt whether a player has sustained a concussion, they should be removed from play and undergo medical assessment by a medical doctor or nurse practitioner.

REFER — Once removed from play, the player with suspected concussion must be referred to a medical doctor or nurse practitioner with training in the evaluation and management of head injury and concussions. All cases of suspected concussion require referral to medical doctors or nurse practitioners for diagnosis, even if the symptoms resolve. In geographic regions of Canada with limited access to medical doctors (rural or northern communities), a licensed healthcare professional (i.e. nurse) with support from a medical doctor or nurse practitioner can provide this diagnostic evaluation.

REPORT — Communication between players, parents, team staff, and their health care providers is vital for the welfare of the player.

Players, parents, and guardians must disclose the nature of, and status of all active injuries (including concussions) to coaches and team staff. Players need to be responsible for one another and encourage the disclosure of concussion symptoms. For children and adolescents with



Chilliwack FC Concussion Management Policy

suspected concussion who have not been directly transferred for medical management, coaches must communicate their concerns directly with the parents or guardians.

RECOVER — Avoiding physical and brain activities that make concussive symptoms worse is the cornerstone of current concussion management. The management of a concussion involves an initial limited period (<24-48 hours) of physical and brain rest. **Stage 1** of the Return-to-Soccer Strategy (see Page 8) involves avoiding or limiting physical and brain activities that make concussive symptoms worse. Once concussion related symptoms have resolved, the player may start **Stage 2** and continue to proceed to the next level when he/she completes the stage without a recurrence of concussion-related symptoms. In conjunction with your school and educational professionals and health care provider, recommendations will be made about whether it is appropriate to take time away from school, or whether returning to school should be done in a graded fashion, this is called “return to learn”. Your health care provider will also make recommendations about whether it is appropriate to take time away from work, or whether returning to work should be done in a graded fashion, this is called “return to work”.

RETURN TO PLAY Players who have been removed from play and referred for medical assessment for a suspected concussion who provide a completed Assessment Medical Report that is signed by a medical doctor or nurse practitioner which documents NO active concussion and may participate in training sessions and game play. Players who have been removed from play and referred for assessment for a suspected concussion who provide a completed Assessment Medical Report that is signed by a medical doctor or nurse practitioner which documents a concussion diagnosis may participate in training sessions (**Stage 3 and 4**) within the Return-to-Soccer Strategy, once they or their parents/guardians report NO concussion symptoms and successfully completing **Stage 2** (15 minutes of light aerobic activity). Players who have concluded **Stage 4** within a Return-to-Soccer Strategy who provide a second completed Assessment Medical Report that is signed by a medical doctor or nurse practitioner which documents recovered concussion may participate in full contact training sessions (Stage 5) and subsequently, game play within the Return-to-Soccer Strategy, if they remain clear of concussion symptoms.

RETURN TO SOCCER STRATEGY Depending on the severity and type of the symptoms, players may progress through the following stages at different rates. Stages 2-4 should each take a minimum of 24 hours in adults, and longer in those 18 years and under.

If the player experiences new symptoms or worsening symptoms at any stage, they should go back to the previous stage and attempt to progress again after being free of concussion-related symptoms for 24 hour or seek medical attention.



Chilliwack FC Concussion Management Policy

Return to Soccer Strategy

	EXERCISE ALLOWED	% MAX HEART RATE	DURATION	OBJECTIVE
STAGE 0 REST	<ul style="list-style-type: none"> ▶ Rest ▶ NO activities 	No training	<1-2 days	Rest
STAGE 1 SYMPTOM LIMITED	<ul style="list-style-type: none"> ▶ Daily activities that do not provoke symptoms. 		Until concussion symptoms clear	Recovery Symptom free
STAGE 2 LIGHT EXERCISE	<ul style="list-style-type: none"> ▶ Walking, light jogging, swimming or stationary cycling at slow to medium pace. ▶ NO soccer ▶ NO resistance training, weight lifting, jumping or hard running 	<70%	<15 min	Increase heart rate.
STAGE 3 SOCCER-SPECIFIC EXERCISE	<ul style="list-style-type: none"> ▶ Simple movement (ie running drills) ▶ Limiting body and head movements. ▶ NO head impact activities. ▶ NO heading. 	<80%	<45 min	Add movement.
STAGE 4 NON-CONTACT TRAINING	<ul style="list-style-type: none"> ▶ Progression to more complex training activities with increased intensity ▶ Coordination and attention e.g. passing, change of direction, shooting, small-sided game ▶ May start resistance training ▶ NO head impact activities ▶ NO heading ▶ Goalkeeping activities should avoid diving and any risk of the head being hit by a ball 	<90%	<60 min	Exercise, coordination, skills/ tactics.
	Youth (<18 years) and adult student-athletes have returned to full-time school activities at this time			
	Repeat medical assessment with second Concussion Assessment Medical Report			
STAGE 5 FULL CONTACT PRACTICE	<ul style="list-style-type: none"> ▶ Normal training activities ie tackling, heading diving saves 	<100%		Restore confidence and assess functional skills by coaching staff.
STAGE 6 GAME PLAY	<ul style="list-style-type: none"> ▶ Normal game play. 	<100%		Player rehabilitated

REASSESS— A player with prolonged concussion recovery (>4 weeks for youth athletes, >2 weeks for adult athletes), or recurrent or complicated concussions, should be assessed and managed by a medical doctor with experience in sports-related concussions, working within a multidisciplinary team.